



(e) If the Lagrangian of a particle moving in one-dimension is given by  $L = \frac{\dot{x}^2}{2x} - V(x)$ , the Hamiltonian is

(i)  $\frac{1}{2}x\dot{p}^2 + V(x)$

(ii)  $\frac{1}{2}p\dot{x}^2 + V(x)$

(iii)  $\frac{1}{2}\dot{x}^2 + V(x)$

(iv)  $\frac{1}{2x}p^2 + V(x)$

(f) Which of the following pairs are canonical?

(i)  $Q = \frac{1}{p}, P = qp^2$

(ii)  $Q = q \cos \beta p, P = q^2 \sin \beta p$

(iii)  $Q = \log(\sin p), P = q \cot p$

(iv)  $P = \frac{1}{2}(p+q), Q = \tan^{-1} \frac{q}{p}$

(g) The equation of motion of system described by the time-dependent Lagrangian

$$L = e^{\gamma t} \left[ \frac{1}{2} m \dot{x}^2 - V(x) \right] \text{ is}$$

(i)  $m\ddot{x} + \gamma m\dot{x} + \frac{dV}{dx} = 0$

(ii)  $m\ddot{x} + \gamma m\dot{x} - \frac{dV}{dx} = 0$

(iii)  $m\ddot{x} - \gamma m\dot{x} + \frac{dV}{dx} = 0$

(iv)  $m\ddot{x} + \frac{dV}{dx} = 0$

(h) If  $q_k, p_k$  are the generalised coordinate and corresponding momentum then the correct relation for Poisson bracket is

(i)  $[q_k, q_l] = \delta_{kl}$

(ii)  $[q_k, q_k] = 1$

(iii)  $[q_k, p_k] = 0$

(iv)  $[q_k, p_k] = 1$

(i) The Routhian for the Lagrangian  $L = \dot{\phi}^2 + r^2 \dot{\theta}^2 + \frac{\mu}{r}$  is given by

(i)  $\frac{p_\theta^2}{4r^2} + \dot{\phi}^2 + \frac{\mu}{r}$

(ii)  $\frac{p_\theta^2}{4r^2} - \dot{\phi}^2 + \frac{\mu}{r}$

(iii)  $\frac{p_\theta^2}{4r^2} - \dot{\phi}^2 - \frac{\mu}{r}$

(iv)  $\frac{p_\theta^2}{4r^2} + \dot{\phi}^2 - \frac{\mu}{r}$

- (j) A dynamical system having kinetic energy  $T$  and potential energy  $V$  is described by the Hamiltonian  $H$ . Assume that the equations defining the generalised coordinates do not depend on time. Then
- (i)  $H = T + V$  and is conserved      (ii)  $H = T + V$  but it is not conserved  
 (iii)  $H \neq T + V$       (iv)  $L = T + V$ .

**Group - B****Unit - 1****(Marks : 10)**2. Answer *any two* questions :

- (a) Define generalised coordinates and a scleronomic system. Find the expression for the kinetic energy of system of  $N$  particles and show that for a scleronomic system the kinetic energy is a homogeneous function of generalised velocities. 2+2+1
- (b) A bead of mass  $m$  slides on a smooth uniform circular wire of radius  $r$  which is rotating with a constant angular velocity  $\omega$  about a fixed vertical diameter. Set up the Lagrangian and find the equation of motion of the bead. 3+2
- (c) A particle of mass  $m$  moving in a central orbit under inverse square law. Construct the Lagrangian and hence find the equation of motion. 2+3
- (d) Define cyclic coordinate. Show that the generalised momentum associated with an cyclic coordinate is a constant of motion for a conservative system. Hence, show that for the motion of a particle in a central force field of potential  $V(r)$ , angular momentum is conserved. 1+2+2

**Unit - 2****(Marks : 15)**3. Answer *any three* questions :

- (a) (i) A particle in two dimensions is in a potential  $V(x, y) = x + 2y$ . Write down the expressions for the canonical momentum  $p_x$  and  $p_y$ . Hence show that  $(p_y - 2p_x)$  is a constant of motion.  
 (ii) The Hamiltonian of a dynamical system of two degrees of freedom is given by

$$H = q_1 p_1 - q_2 p_2 - a q_1^2 + b q_2^2,$$

where  $a$  and  $b$  are constants. Show that  $F_1 = \frac{p_1 - a q_1}{q_2}$  is a constant of motion. 3+2

- (b) A particle with coordinate  $q$  and momenta  $p$  has the Hamiltonian  $H = qp^2 - qp + bp$ , where  $b$  is a constant. Solve for  $p$  and  $q$ . 3+2
- (c) A particle of mass  $m$  is attracted towards a given point by a central force of the form  $k/r^2$ , where  $k$  is a constant. Assuming that the central force is conservative in nature, write down the expression for the Hamiltonian of the system and derive Hamilton's equations of motion. 3+2

**Please Turn Over**

- (d) If the Lagrangian  $L_0 = \frac{1}{2}m\left(\frac{dq}{dt}\right)^2 - \frac{1}{2}m\omega^2 q^2$  of a dynamical system is modified to  $L = L_0 + \alpha q\left(\frac{dq}{dt}\right)$ , then show that canonical momentum changes though equation of motion does not change. 2+3
- (e) State Hamilton's principle. Derive Hamilton's principle from D'Alembert's principle.

### Unit - 3

(Marks : 10)

4. Answer *any two* questions :

- (a) A particle of mass  $m$  moves on the inside of a frictionless cone having equation  $x^2 + y^2 = z^2 \tan^2 \alpha$ . Using cylindrical coordinates write the Hamiltonian and Hamilton's equations. 3+2
- (b) Show that the integral

$$J_1 = \iint_S \sum_i dq_i dp_i$$

taken over an arbitrary two-dimensional surface  $S$  of the  $2n$ -dimensional phase space is invariant under canonical transformation. 5

- (c) (i) Using Hamilton's equations of motion, show that the Hamiltonian  $H = \frac{p^2}{2m} e^{-rt} + \frac{1}{2} m \omega^2 x^2 e^{rt}$  leads to the equation of motion of a damped harmonic oscillator  $\ddot{x} + r\dot{x} + \omega^2 x = 0$ .
- (ii) If the Lagrangian of a two-dimensional Harmonic oscillator is  $L = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\theta}^2) - \frac{1}{2}kr^2$ , where  $k$  is a constant, then determine the Hamiltonian of the system. 3+2
- (d) A system is governed by the Hamiltonian

$$H = \frac{1}{2}(p_x - ay)^2 + \frac{1}{2}(p_y - bx)^2,$$

where  $a, b$  are constants and  $p_x, p_y$  are momenta conjugate to  $x$  and  $y$  respectively. Find the values of  $a$  and  $b$  so that the quantities  $(p_x - 3y)$  and  $(p_y + 2x)$  are conserved. 5

(5)

Z(6th Sm.)-Mathematics-H/DSE-B-2.3/CBCS

Unit - 4

(Marks : 10)

5. Answer *any two* questions :

- (a) When is a transformation said to be a canonical transformation? A canonical transformation  $(q, p) \rightarrow (Q, P)$  is made through the generating function  $F(q, p) = q^2 p$  on the Hamiltonian

$$H(q, p) = \frac{p^2}{2\alpha q^2} + \frac{\beta}{4} q^4, \text{ where } \alpha, \beta \text{ are constants. Find the equation of motion for } (Q, P).$$

1+4

- (b) Show that the transformation  $q = \sqrt{2P} \sin Q$  and  $p = \sqrt{2P} \cos Q$  is canonical. Find a generating function for this transformation. 5
- (c) If the Poisson bracket of a time-independent dynamical variable  $u$  with the Hamiltonian concerned vanishes, show that  $u$  is a constant of motion. 5
- (d) State and prove Liouville's theorem for a Hamiltonian system. 5
-